

Chapter 8

1. A 5.00-kg block is set into motion up an inclined plane with an initial speed of $v_i = 8.00$ m/s (Fig. P8.23). The block comes to rest after traveling $d = 3.00$ m along the plane, which is inclined at an angle of $\theta = 30.0^\circ$ to the horizontal. For this motion, determine (a) the change in the block's kinetic energy, (b) the change in the potential energy of the block-Earth system, and (c) the friction force exerted on the block (assumed to be constant). (d) What is the coefficient of kinetic friction?

Ans:




Figure P8.23

2. An older-model car accelerates from 0 to speed v in a time interval of Δt . A newer, more powerful sports car accelerates from 0 to $2v$ in the same time period.

Assuming the energy coming from the engine appears only as kinetic energy of the cars, compare the power of the two cars.

Ans:

Chapter 9.

1. An object has a kinetic energy of 275 J and a momentum of magnitude $25.0 \text{ kg} \cdot \text{m/s}$.

Find the speed and mass of the object.

Ans:

2. When you jump straight up as high as you can, what is the order of magnitude of the maximum recoil speed that you give to the Earth? Model the Earth as a perfectly solid object. In your solution, state the physical quantities you take as data and the values you measure or estimate for them.

Ans

:

3. The front 1.20 m of a 1 400-kg car is designed as a “crumple zone” that collapses to absorb the shock of a collision. If a car traveling 25.0 m/s stops uniformly in 1.20 m, (a) how long does the collision last, (b) what is the magnitude of the average force on the car, and (c) what is the acceleration of the car? Express the acceleration as a multiple of the acceleration due to gravity.

Ans:

4. A railroad car of mass 2.50×10^4 kg is moving with a speed of 4.00 m/s. It collides and couples with three other coupled railroad cars, each of the same mass as the single car and moving in the same direction with an initial speed of 2.00 m/s. (a) What is the speed of the four cars after the collision? (b) How much mechanical energy is lost in the collision?

Ans:

5. Four objects are situated along the y axis as follows: a 2.00-kg object is at +3.00 m, a 3.00-kg object is at +2.50 m, a 2.50-kg object is at the origin, and a 4.00-kg object is at -0.500 m. Where is the center of mass of these objects?

Ans: